



Burnt Bridge Creek

Summary of 2018 Surface Water Monitoring Program Results | November 2019



The Washington State Department of Agriculture (WSDA) routinely monitors surface water throughout the state for the presence of pesticides. The monitoring is done between March and September, the typical season for pesticide use, and includes checking general water quality conditions and streamflow. State and federal agencies use this data to evaluate water quality and make exposure assessments for pesticides registered for use in Washington State. In 2018, WSDA monitored 16 sites in Washington, one of them in Clark County.



Watershed and site information

Sampling history: 2017 - Present

Watershed area: 16,800 acres (~26 square miles)

Area in agricultural use: 500 acres (~3% of total watershed acreage)

Main crops: Golf course, grass hay, CRP/conservation, and pasture

Fish habitat: Winter steelhead and coho salmon (SalmonScape: apps.wdfw.wa.gov/salmonscape/)

Sampling dates: 14 sampling visits, March 27 – Sept. 25, once every 2 weeks

Water testing:

- WSDA tested for 144 current and legacy chemicals (50 insecticides, 54 herbicides, 20 fungicides, 15 pesticide degradates, 2 synergists, 1 antimicrobial, 1 insect repellent, and 1 wood preservative).
- Samples were analyzed at Manchester Environmental Lab, Port Orchard, Washington.
- WSDA compares detected pesticide concentrations to WSDA assessment criteria, which are half of state and federal water quality criteria. Each pesticide has its own assessment criteria, based on its toxicity to aquatic animals, insects, and plants.
- WSDA identifies Pesticides of Concern (POCs) as current-use pesticides that have been found somewhere in the state above WSDA's assessment criteria in recent years.

Notes:

- The creek flows through 10 miles of Vancouver, Washington's residential and agricultural areas.
- Historical data suggests Burnt Bridge is the least healthy stream in Clark County.
- Stream habitat has improved because of efforts by non-profits, volunteers, and government agencies. Their work included riparian vegetation plantings and stormwater drainage control.

Results and Conclusions

- There were 218 total pesticide detections in Burnt Bridge Creek from 5 different use categories: 21 types of herbicides, 6 fungicides, 5 insecticides, 7 degradates, and 2 other pesticide-related chemicals. This substantial increase from 2017 is largely due to new equipment at the lab and does not necessarily reflect an increase in pesticide use.
- Of the total pesticide detections, 14 were above WSDA's assessment criteria.
- The POCs diuron, fipronil, metolachlor, pyridaben, pyriproxyfen, and cis-permethrin were detected.
- The only POC detected at this site at a concentration that exceeded WSDA's assessment criteria was cis-permethrin.
- When multiple pesticides are detected simultaneously, the environmental effects can combine; multiple pesticides were detected every week Burnt Bridge Creek was tested. Between 7 and 21 pesticides were detected at each sampling visit.

Recommendations


- **Make use of natural protections**
 - Use buffers, filter strips, sediment basins, ground cover, and setbacks.
- **Be informed**
 - Read and follow pesticide label directions, and be familiar with active ingredients.
 - Plan applications using the weather forecast to reduce the chances of drift or runoff.
 - Review WSDA's POCs and choose less-toxic pesticides when possible.
- **Care for your equipment and products**
 - Calibrate, maintain, and inspect application equipment regularly.
 - Properly dispose of all unneeded pesticides. Visit agr.wa.gov/wastepesticide to learn about waste pesticide collection events.

The calendar below shows the concentration in µg/L and date sampled of each WSDA POC. This calendar does not include all the pesticides WSDA found during the growing season. Detected concentrations that exceed WSDA's assessment criteria have a higher potential to cause harm to aquatic ecosystems. The “-” signifies a sample or measurement that was not collected or could not be analyzed.

Washington State's Pesticides of Concern Detected and their Corresponding Sampling Dates and Concentrations														
Month	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sept.							
Day of the Month	Use*	11	25	9	23	6	20	2	18	1	15	29	12	25
Diuron	H	0.004	0.003	0.003	0.005								0.006	
Fipronil	I				0.003									
Metolachlor	H	0.003										0.002	0.003	0.002
Pyridaben	I													0.004
Pyriproxyfen	I										0.004			0.004
cis-Permethrin	I													0.006
Total suspended solids (mg/L)	13.0	12.0	8.0	12.0	12.0	6.0	-	7.0	6.0	6.0	5.0	4.0	13.0	3.0
Streamflow (cubic ft/sec)	25.53	24.95	14.84	11.16	9.11	7.44	7.45	6.41	4.70	4.00	4.44	4.89	5.82	4.76
Precipitation (total in/week)	1.53	1.25	0	0.01	0	0.09	0.16	0	0	0	0	0.07	0.12	0

Exceeds Assessment Criteria
 Below Assessment Criteria
 (* H: Herbicide, I: Insecticide)

The graph below shows the total number of detections per sampling visit in each pesticide category. The category 'other' includes degradates and additional pesticide-related chemicals. Note that the number of detections between categories cannot be directly compared due to the different number of chemicals in each category and variability in analysis methods used. Bars without data labels represent one detection for that sampling visit and pesticide category.



In the triangle to the right, pesticides in the top section have one or more detections above WSDA assessment criteria. The total number of detections for each pesticide is in parentheses after the name, with more frequently detected pesticides listed first in each section.

Please see agr.wa.gov/AgScience for more information.

