


FINAL
ENVIRONMENTAL ASSESSMENT
AND
FINDING OF NO SIGNIFICANT IMPACT

COOPERATIVE GYPSY MOTH ERADICATION PROJECT
PIERCE COUNTY
WASHINGTON

April 17, 2012



Prepared by
Washington State Department of Agriculture
Plant Protection Division

In cooperation with
United States Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA'S TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 1400 Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

Mention of companies or commercial products in this report does not imply recommendation or endorsement by USDA over others not mentioned. USDA neither guarantees nor warrants the standard of any product mentioned. Product names are mentioned solely to report factually on available data and to provide specific information.

This publication reports research involving pesticides. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.

TABLE OF CONTENTS

| | |
|---|----|
| I. PURPOSE AND NEED FOR ACTION..... | 1 |
| A. Decisions to be Made and Scope of Analysis..... | 1 |
| 1. Introduction..... | 1 |
| 2. Environmental Analysis and Documentation..... | 1 |
| 3. History and Scope of Project..... | 2 |
| 4. Decisions to be Made..... | 3 |
| B. Need For Action..... | 3 |
| 1. Project Goals and Objectives..... | 4 |
| C. Proposed Action..... | 4 |
| D. Authorizing Laws and/or Policies..... | 4 |
| 1. State Authorizing Laws..... | 4 |
| 2. Federal Authorizing Laws..... | 4 |
| 3. Environmental Laws and Other Regulations..... | 5 |
| II. PUBLIC INVOLVEMENT AND ISSUES..... | 5 |
| A. Public Notification and Involvement..... | 5 |
| B. Issues and Concerns..... | 7 |
| III. AFFECTED ENVIRONMENT..... | 8 |
| A. 2011 Site Descriptions..... | 8 |
| B. Threatened, Endangered, and Sensitive Species..... | 10 |
| IV. TREATMENT ALTERNATIVES..... | 11 |
| A. Treatment Alternatives..... | 11 |
| B. Preferred Treatment Alternative..... | 12 |
| C. Treatment Alternatives Not Selected..... | 12 |
| V. ENVIRONMENTAL CONSEQUENCES..... | 13 |

| | |
|---|----|
| A. Human Health and Safety..... | 13 |
| 1. <u>No Action</u> | 13 |
| 2. Proposed Action | 14 |
| B. Non-Target Organisms | 16 |
| 1. Bacillus thuringiensis var. (kurstaki) (Btk.)..... | 16 |
| 2. Bond Max | 17 |
| VI. MONITORING | 17 |
| VII. CUMULATIVE EFFECTS | 17 |
| VIII. SUMMARY | 18 |
| IX. LIST OF AGENCIES AND PERSONS CONSULTED | 19 |
| X. LIST OF PREPARERS..... | 20 |
| XI. APPENDICES | 20 |
| A. References | |
| B. Treatment Site Maps | |
| C. Standard Operating Procedures | |
| D. Product Labels and Material Safety Data Sheets | |
| E. Finding of No Significant Impact (FONSI) | |

I. PURPOSE AND NEED FOR ACTION

A. Decisions to be Made and Scope of Analysis

1. Introduction

The Washington State Department of Agriculture (WSDA), in cooperation with the United States Department of Agriculture Animal and Plant Health Inspection Service (USDA-APHIS), is proposing an eradication program with the goal of eliminating two isolated infestations of the non-native gypsy moth, Lymantria dispar (Linnaeus), in Pierce County, Washington in the spring of 2012.

2. Environmental Analysis and Documentation

In 1995, the USDA Forest Service and APHIS prepared a final environmental impact statement, "Gypsy Moth Management in the United States: a cooperative approach", (hereinafter referred to as FEIS), which described and analyzed methods of gypsy moth control available for use in USDA cooperative programs.

This Environmental Assessment (EA) is tiered to the FEIS in accordance with the Council on Environmental Quality regulations for implementing the National Environmental Policy Act of 1969 (NEPA) (40 CFR 1502.20 and 40 CFR 1508.28). This EA provides the basic background information necessary for the site-specific analysis of the potential environmental effects of WSDA's proposed 2012 Cooperative Gypsy Moth Eradication Project. The FEIS and this site-specific EA jointly constitute the environmental analysis and documentation required under NEPA.

Copies of the FEIS and the EA are available for review at:

Washington State Library
6880 Capitol Blvd. S
Tumwater, WA 98501

and

USDA, APHIS, PPQ
APHIS Library, 1st floor
4700 River Road
Riverdale, MD 20737

and

USDA, APHIS, PPQ
33400 9th Ave. S., Suite 200
Federal Way, WA 98003

Additional environmental analysis and documentation has been prepared to satisfy Washington State requirements under Chapter 43.21 (c) of the Revised Code of Washington (State Environmental Policy Act or SEPA), and Chapter 197-11 of the Washington Administrative Code (SEPA rules).

Copies of the SEPA documentation are available for review at:

Washington State Library
6880 Capitol Blvd. S
Tumwater, WA 98501

and

Washington State Department of Agriculture
www.agr.wa.gov

3. History and Scope of Project

Since its accidental release in the United States in 1869, the European strain of gypsy moth has spread throughout New England and areas to the north, south and west. It has become established in all or parts of 19 states, the District of Columbia, and parts of Canada. It continues to spread to uninfested areas. The gypsy moth has caused dramatic economic, social, and ecological impacts throughout the infested area (USDA, 1995, vol. II, chapter 1, p. 4).

The European strain of the gypsy moth has been found every year in Washington State since 1974 with the exceptions of 1976 and 1977. The European gypsy moth is usually introduced to Washington State by people visiting or relocating from the infested area of eastern North America. For more than 30 years, WSDA has successfully detected new introductions of European strain of gypsy moth and successfully eradicated all reproducing populations.

In 1991, the Asian strain of the gypsy moth was found for the first time in Oregon, Washington, and in British Columbia, Canada. Eradication projects conducted in 1992 successfully eliminated the insect from those areas. WSDA has detected and treated introductions of the Asian strain of the gypsy moth in 1991-92, 1994-95, 1995-96, 1996-97, 1997-98 and 1999-2000. These eradication projects have been successful. The Asian strain poses a far greater risk of rapid spread than the European. Unlike females of the European strain, females of the Asian strain may fly and deposit an egg mass miles from where they feed as caterpillars. The Asian strain also poses a greater risk of damage because it feeds on a greater variety of plants (USDA, 1995, vol. II, chapter 1, p. 4).

For more information on how the different strains/populations of the gypsy moth are to be treated please see USDA, 1995, vol. II, chapter 1, pp. 9-11.

4. Decisions to be made

There are three significant decisions which must be made as a part of evaluating a gypsy moth control action.

The first decision to be made is whether to propose a gypsy moth control project (the absence of a control project is a no-action alternative). The second decision to be made is whether or not tiering this environmental assessment to the USDA 1995 FEIS is appropriate. The third decision to be made is what tools are being proposed for the project area.

B. Need For Action

In order to avoid undesirable economic, social, and ecological impacts to residents, communities and businesses in Washington State, WSDA in cooperation with USDA APHIS, proposes to eradicate two isolated infestations of European gypsy moth in Pierce County. One proposed treatment area is in the city of **Puyallup**.

Seven adult male gypsy moths were caught in Puyallup during WSDA's summer trapping program in 2010. Follow-up inspections in the area of the catches revealed alternate gypsy moth life stages (pupal cases and egg masses) indicating the existence of a reproducing population. WSDA conducted a ground treatment in Puyallup (29 acres) in spring of 2011. Subsequent surveys in summer and fall of 2011 led to the discovery of 10 male moths (trapped), 3 larvae, 2 pupal cases, and one viable egg mass in or near the treatment area. The lack of success of the 2011 Puyallup treatment is likely due to an extremely late spring (and subsequent late egg hatch) as evidenced by some very late (Sept. 22) live adult moths trapped in 2011.

The second proposed treatment area is in the city of **Eatonville**.

Two adult male gypsy moths were caught in Eatonville during the 2011 summer trapping program. Follow up inspections of the area revealed a resident had recently moved into the neighborhood from a gypsy moth infested state (Michigan). Further inspections revealed a bird house had been transported from Michigan containing viable egg masses (spent egg masses found in the bird house). A viable egg mass was then discovered near the property.

Evidence of an isolated reproducing population of European gypsy moth in Washington is a "trigger" to evaluate eradication options.

1. Project Goals and Objectives

The WSDA, in cooperation with USDA-APHIS and other appropriate Federal, State and local agencies, proposes to take action to eradicate two isolated infestations of European gypsy moth in Pierce County, Washington. The action will be designed to give the project the best chance for achieving the goal of eradicating the gypsy moth infestation while minimizing risks to human health as well as minimizing detrimental environmental consequences. This action will be taken in order to prevent the establishment and spread of this pest insect and thereby avoid the adverse economic, social, and ecological effects associated with large-scale gypsy moth infestations.

C. Proposed Action

Strategies described in the FEIS depend upon the infestation status of the area: generally infested, transition, or uninfested. The three strategies of suppression, eradication, and slow the spread -- or their absence -- are included in the six alternatives described in the FEIS. The sixth alternative is the preferred alternative presented in the FEIS. The sixth alternative is comprised of all three strategies.

Based on the infestation status of "no established population", Washington State's strategy in 2012 will be eradication.

Treatments available for eradication projects include: (the biological insecticides) Bacillus thuringiensis var. kurstaki (Btk) and the gypsy moth nucleopolyhedrosis virus (Gypchek); a chemical insecticide (diflubenzuron); and treatments employing mass trapping, mating disruption, and sterile insect release techniques. A detailed description of these treatment options and the decision making process can be found in Section IV of this EA.

After evaluating treatment options available in the 1995 FEIS, WSDA proposes three to five applications of Btk to 43 acres of vegetation at the core of the infestation in Puyallup and to 13 acres at the core of the infestation in Eatonville. The applications will target early instar larvae shortly after egg hatch in April and May.

D. Authorizing Laws and/or Policies

1. State Authorizing Laws

WSDA has authority under Chapter 17.24 of the Revised Code of Washington, Insect Pests and Plant Diseases, to eradicate or control insect pests that may endanger the agricultural and horticultural industries in the state of Washington.

2. Federal Authorizing Laws

The USDA-APHIS has broad discretionary authority to prevent the establishment or spread of plant pests. See 1995 FEIS, volume 2, chapter 1, pages 8 and 9, "Statutory Authorities", for more information.

3. Environmental Laws and Other Regulations

Many environmental laws, authorities and Executive Orders of the President influence how actions to manage pests, including the gypsy moth, are implemented at the site-specific level. Such laws include the National Environmental Policy Act; the Washington State Environmental Policy Act; the Federal Insecticide, Fungicide, and Rodenticide Act; the Clean Water Act and the Endangered Species Act. See 1995 FEIS, volume 2, chapter 1, pages 8 and 9, "Statutory Authorities", for more information.

II. PUBLIC INVOLVEMENT AND ISSUES

A. Public Notification and Involvement

WSDA conducts on-the-ground inspections in early fall 2011. Washington State Department of Agriculture (WSDA) employees searched for egg masses and other evidence of gypsy moth activity in numerous communities (including Puyallup and Eatonville) where multiple moth catches had been made in summer 2011. During these inspections, WSDA representatives had the opportunity to interact with many local residents and to explain the purpose and value of WSDA's gypsy moth program.

WSDA delivers letters to locally elected officials in Puyallup and Eatonville on December 20, 2011. Officials receiving letters included the state senator and two state representatives from the 2nd and the 25th legislative district, members of the Pierce County Council, and mayor and city council members of Puyallup (and City Manager) and Eatonville. The letters stated:

1. A reproducing population of gypsy moth had been located in the cities of Puyallup and Eatonville.
2. WSDA is proposing to eradicate the infestation with a biological insecticide *Bacillus thuringiensis* var. *kurstaki* (*Btk*) in spring 2012.
3. WSDA will soon begin a public information campaign to inform local residents and community leaders of the infestation and proposed treatment.

WSDA delivers letters to residents and businesses in or near the proposed treatment zones on December 21, 2012. The letters stated:

1. A reproducing population of gypsy moth exists in your neighborhood.
2. WSDA is proposing a series of treatments of a biological insecticide, *Bacillus thuringiensis* var. *kurstaki*, beginning in April or May to eradicate the destructive pest.

3. You are invited to an open house (details were contained in the letter) to learn more about the proposed treatment.
4. Please call WSDA's toll-free hotline (1-800-443-6684) or visit the WSDA web site at www.agr.wa.gov, click on gypsy moth, for more information.

Enclosed with the letter were a gypsy moth fact sheet and a map of the proposed treatment site.

WSDA dispatches news release to local media December 27, 2011: The news release announced the proposal to treat a 43-acre site at the South Hill Mall in the city of Puyallup and a 13-acre site in the city of Eatonville in the spring of 2012. Also included in the news release was the time frame of treatments, WSDA's compliance with environmental review policy, proposed insecticide to be used, and announcement of community open houses to take place in early February. Readers were encouraged to call the agency's toll-free hotline or visit the WSDA gypsy moth website for more information.

WSDA dispatches news release to local media January 25, 2012: The news release announced the details of two public open houses to be held in February of 2012. The Eatonville open house is scheduled for February 7th and the Puyallup open house is scheduled for February 13th.

WSDA dispatches an email to stakeholders and local elected officials on February 3, 2012: The email stated that a community open house would be held in Eatonville on February 7th and in Puyallup on February 13th to:

1. Discuss strategies and treatments for addressing gypsy moth infestation in Washington.
2. Discuss why eradication is the strategy selected to respond to infestations in Washington.
3. Discuss the process used by WSDA to evaluate and propose a treatment.
4. Inform the public of the opportunity to comment on the SEPA and NEPA documents.

WSDA holds community open house in Eatonville on February 7, 2012 and in Puyallup on February 13. The open houses were held at Eatonville High School and Woodland Elementary School (Puyallup). Subject matter experts from WSDA and Tacoma/Pierce County Health Department were present to provide information and answer questions. Attendees were able to visit five different stations at the open house: 1) Trapping data; 2) Proposed treatment zone; 3) Technical reference table; 4) Human Health issues; 5) 12-minute DVD presentation

WSDA emphasized several major points at open houses:

1. Destructiveness of the gypsy moth.
2. How the pest gets to Washington.
3. How the pest damages the environment and the economy.
4. Results of WSDA's summer trapping program.
5. Evidence supporting the eradication proposal.
6. Boundaries of the proposed treatment zone.

7. Proven safety record of the pesticide proposed for use.
8. The two environmental documents made available for public review and comment for an eradication proposal.
9. The opportunity residents have to comment on the environmental documents.
10. Treatments available to control gypsy moths.
11. Why eradication is the best strategy for Washington.
12. Failure of early attempts in the late 1800s, 1900s to eradicate the moth.
13. Where 85 treatments have been conducted in Washington.
14. The process WSDA follows when deciding whether or not to conduct a treatment.

Attendees also received a packet to take home with them containing the following handouts:

1. Why the gypsy moth is one of the worst pests ever brought into the U.S.
2. How the gypsy moth damages the environment
3. Purpose of gypsy moth open houses
4. Background data on the pesticide commonly used in eradication treatments
5. Washington State Department of Health fact sheet
6. Map of the proposed treatment zone
7. Map showing the spread of the gypsy moth in U.S. from 1900 to 2000
8. Photos of America's first major gypsy moth outbreak in 1889
9. Where 85 gypsy moth eradication treatments have been conducted in Washington since 1979
10. Advantages and disadvantages of six treatments available to WSDA to control gypsy moths
11. The eight steps WSDA goes through in deciding to conduct an eradication treatment
12. Why eradication is the best of four basic strategies for Washington

Environmental review documents available for public comment. This EA and the State Environmental Policy Act (SEPA) review documents will be made available for a 30-day public comment period. Notice of availability will be advertised in local and regional newspapers. Documents will be distributed to stakeholders, made available at local libraries, and posted on the WSDA and USDA websites.

B. Issues and Concerns

Among the attendees at the open houses was the Mayor of Eatonville, a city council member from the city of Eatonville, and several local residents.

- Among the questions attendees asked at the open house were the following:

Q: "When will the treatments be administered?" (Answer: April/May time frame)

Q: "What kind of damage does the gypsy moth do?" (Answer: The gypsy moth causes extensive defoliation. The person was then shown photos of damaged vegetation and encouraged to watch the 12-minute DVD.)

Q: "Are there any ill effects from the insecticide treatment?" (Answer: Btk is not considered a public health risk. In the past a handful of people have reported mild skin reactions or mild eye, ear, and nose irritations after Btk treatments have been administered. However, health officials have never been able to determine if the reactions were caused by Btk; pollens, molds, or dust generated by the treatment; or were unrelated to the treatment.)

Q: "Has there been any opposition to the treatment?" (Answer: None to date. WSDA will continue to conduct an aggressive public information campaign, to ensure the public receives timely, accurate information and supports our proposal)

Q: "We strongly support eradication treatments. Can we submit a written statement of support?" (Answer: Yes. WSDA is very happy to receive statements of support for eradication proposals.)

Q: "Will you spray if it's raining?" (Answer: No. If steady rain is falling or forecast, we will postpone the treatment and wait for more favorable weather.)

III. AFFECTED ENVIRONMENT

A. 2011 SITE DESCRIPTION (see Appendix B for maps)

Puyallup (Puyallup, WA 7.5 minute quadrangle, S4 T19N R4E)

- Pierce County, Washington
- Approximately 43 acres
- Zoning
GC: General Commercial
- Proposed Area
The proposed 43 acre site is primarily located in the parking area of the South Hill Mall. (See site map in Appendix B)
- Vegetation
The site is primarily a paved parking area. Vegetation consists of landscape trees and shrubs and some volunteer trees along the north boundary fence line.

Canopy coverage is less than 10%, tree height is variable with deciduous trees in excess of 50 feet.

- **Critical/Sensitive Areas**

There is a small wetland area (0.7 acres) along the southern boundary of the treatment area. WSDA will be working under NPDES permit #WA0039047 issued by the Department of Ecology to WSDA for the purpose of invasive moth control. The permit, titled "Invasive Moth Control National Pollutant Discharge Elimination System Waste Discharge Permit", authorizes discharge of insecticides into surface waters of the state of Washington that are consistent with the terms and conditions of this permit for the purpose of invasive moth control.

There is one area of steep slope (45%) along the north boundary of the proposed treatment area (See topography map in Appendix B). This steep embankment leads from the highway up to the treatment area. The host vegetation to be treated is primarily at the top of the embankment and will be approached from the top. No disruption or erosion of the embankment is anticipated.

- **Catch History**

Ten European Gypsy Moths were caught in the area during the 2011 summer trapping survey.

- **Alternate Life Stages**

Three GM larvae were found in August 2011.

Two pupal cases were found in the area during the fall of 2011.

One egg mass was found in the area during the fall of 2011.

Eatonville (Eatonville, WA 7.5 minute quadrangle, S15 T16N R4E)

- **Pierce County, Washington**

- **Approximately 13 acres**

- **Zoning**

SF 1: Single Family Residential

MF-1: Multifamily Residential

- **Proposed Area**

The proposed 13 acre site is located in a residential area on the west edge of the City of Eatonville. (See site map in Appendix B)

- **Vegetation**

The site is primarily a paved parking area. Vegetation consists of landscape trees and shrubs and some volunteer trees along the north boundary fence line.

Canopy coverage is approximately 30%, tree height is variable with deciduous trees in excess of 30 feet.

- **Critical/Sensitive Areas**

There is a small pond (0.7 acres) along the southern boundary of the treatment area. WSDA will be working under NPDES permit #WA0039047 issued by the Department of Ecology to WSDA for the purpose of invasive moth control. The permit, titled "Invasive Moth Control National Pollutant Discharge Elimination System Waste Discharge Permit", authorizes discharge of insecticides into surface waters of the state of Washington that are consistent with the terms and conditions of this permit for the purpose of invasive moth control.

- **Catch History**

Two European Gypsy Moths were caught in the area during the 2011 summer trapping survey.

- **Alternate Life Stages**

One egg mass was found in the area during the fall of 2011.

B. Threatened, Endangered, and Sensitive Species

As required by Section 7 of the Endangered Species Act of 1973, the USDA has conferred with both the United States Fish and Wildlife Service (USFWS) and the National Marine Fisheries Service (NMFS). No listed, designated, proposed, or candidate species occur at or near the proposed treatment site. USDA-APHIS has determined that the proposed eradication project will have no effect on any listed, designated, proposed, or candidate species or their critical habitat.

In addition, the WSDA has consulted with the Washington State Department of Fish and Wildlife (WDFW) and the Washington State Department of Natural Resources (DNR). These agencies provided maps or other data intended to aid in the identification of habitats of concern and the presence of listed, proposed, candidate, threatened or endangered species. The information provided by WDFW Priority Habitats and Species Program did not identify any threatened or endangered species on the **Puyallup** or **Eatonville** site. WDFW did identify the following listed species as occurring in the general area:

Three bald eagle nesting sites were listed as occurring in the area of the **Puyallup** treatment site. The listed nesting sites are over one mile from the proposed site

The Nisqually River Corridor (**Eatonville**) is a Bald Eagle use area. Regular sightings occur within five miles of the treatment area.

The Mashel River is approximately 0.5 miles from the **Eatonville** treatment area. Priority fish present in the Mashel include: Coho Salmon, Fall Chinook, Pink Salmon, Resident Cutthroat, Sockeye Salmon, Winter Chum, and Winter Steelhead.

Lynch Creek is approximately 1.0 miles from the **Eatonville** treatment site and is the discharge point for stormwater runoff. Priority fish present in Lynch Creek include: Coho Salmon, Fall Chinook, Pink Salmon, Resident Cutthroat, Sockeye Salmon.

The information provided by WDFW from their lepidopteran database found no butterfly species of concern in the immediate area or within a 5-mile radius of either area.

A review of the DNR Washington Natural Heritage Program database found no records for rare plants or high quality native ecosystems in the vicinity of the **Puyallup** or **Eatonville** site.

IV. TREATMENT ALTERNATIVES

A. Treatment Alternatives

WSDA is proposing to conduct an Integrated Pest Management (IPM) program to eradicate gypsy moth in Washington State. Integrated Pest Management involves selecting those options and techniques that give the best chance of meeting the project goal of eradication. The FEIS contains a range of alternatives from which WSDA has selected an IPM strategy. The treatment alternatives detailed in the FEIS include:

- 1) *Bacillus thuringiensis var kurstaki* (Btk). This is a biological insecticide containing the bacterium Btk. The insecticide is effective primarily against caterpillars of many species of moths and butterflies.
- 2) Diflubenzuron (Dimilin®). This is an insect growth regulator that interferes with the growth of some immature insects.
- 3) Gypsy moth virus (Gypcheck). This is a nucleopolyhedrosis virus which occurs naturally and is specific to GM. Gypcheck is an insecticide product made from the GM nucleopolyhedrosis virus.
- 4) Mass trapping. This treatment consists of large numbers of pheromone traps used to attract the male GM and prevent them from mating with females, thereby causing a population reduction. The density of traps in this treatment option is nine or more traps per acre.
- 5) Mating disruption. This treatment consists of applying tiny plastic flakes or beads containing disparlure, a synthetic GM sex pheromone. The pheromone confuses male moths and, thus, prevents them from locating and mating with females.
- 6) Sterile insect technology. This treatment consists of an aerial release of a large number of sterile male GMs. This reduces the chance that female moths will mate with fertile males. The result is progressively fewer and fewer fertile egg masses being produced, and eventual elimination of the population.

B. Preferred Treatment Alternative

The WSDA/USDA-APHIS gypsy moth eradication project IPM strategy proposed for 2011 includes the use of the biological insecticide (B.t.k). Ground-based equipment will be utilized at the Puyallup site. Ground-based applications may include the spreader-sticker Bond Max. Treatments will also include visual inspections for and removal of egg masses when found, and be followed up by delimitation trapping. This IPM strategy will give the project the best chance to achieve the goal of eradicating the gypsy moth infestations while minimizing risks to human health and minimizing detrimental environmental consequences.

Details of the proposed application:

Ground-based applications will involve three-five treatments of Foray XG (EPA Reg. No. 73049-46) Bacillus thuringiensis var. kurstaki (Btk) applied at label rate. The treatments would occur during the period between April 15 and June 30, 2011. Exact timing of the applications would be dependent on development of gypsy moth larvae and/or foliage as determined by WSDA.

Spreader-sticker (Bond Max) has been common practice for other gypsy moth projects (USDA, 1995, vol. II, A-4). The spreader-sticker ensures that Btk adheres to the foliage rather than falling to the ground.

All ground applications will be conducted in accordance with all applicable federal, state, and local laws and regulations, and will adhere to the Standard Operating Procedures developed by WSDA for this project. See Appendix C.

Follow up:

A follow up trapping program employing pheromone-baited traps in the summer of 2012 will contribute to the success of the eradication project by removing males from any residual population, delimiting the location of any residual populations of Gypsy moths, and aiding in the evaluation of the project.

In the event of multiple moth catches in a treatment area, visual inspections for alternate life stages (egg masses etc.) will be performed in the fall of 2012. Visual inspection will help determine if re-treatment actions should be considered.

C. Treatment Alternatives Not Selected

The other treatment options were not selected due to environmental or efficacy concerns. Diflubenzuron is an insect growth regulator that has adverse impacts on a broader range of nontarget species than Btk. While Btk primarily impacts moths and butterflies, diflubenzuron can kill many other insects in addition to moths and butterfly caterpillars. Its use may adversely affect other insect populations and, therefore, it was not selected. GM

virus (Gypcheck) is very host-specific but is not widely available in the market; it is still somewhat experimental for eradication programs and, therefore, was not selected. Mass trapping has been used with some success to eradicate isolated populations, but at other times has failed. It is best employed following larval pesticide treatments in small, isolated low-level populations. Mating disruption is used primarily in areas to prevent or slow the spread of GM. Sterile insect releases have been approved but have rarely, if ever, been used in eradication efforts.

V. ENVIRONMENTAL CONSEQUENCES

A. Human Health and Safety

1. No Action

The gypsy moth is able to survive and reproduce in Washington State, as evidenced by numerous past isolated infestations. The current infestation, if left unchecked, could spread across a large area.

Trees in forests and orchards, and residential and municipal shade trees and landscape plantings would be damaged and killed. Recreational and aesthetic values associated with trees and forested land would be diminished (USDA, 1995, vol. II, chapter 2, p. 29). Species composition of the vegetation on forested land could change, affecting the quantity and variety of food available for wildlife (USDA, 1995, vol. II, chapter 2, p. 23).

Water quality could be adversely affected in a number of ways including: 1) increased siltation from rapid runoff of rainfall from defoliated areas; 2) increases in water temperature as it flows through areas made shadeless; and 3) nutrient overloading from the deposition of large quantities of caterpillar droppings (USDA, 1995, vol. II, chapter 2, pp. 24-25).

The pesticide load in the environment would likely increase in quantity, variety, and net detrimental environmental impact as home and business owners respond to ever-increasing numbers of gypsy moth caterpillars, the damage they cause, and the nuisance they represent (USDA, 1995, vol. II, chapter 4, p. 76).

Human health effects associated with the presence of large numbers of gypsy moth caterpillars have been reported, including rashes and welts typical of allergic reactions, and respiratory complaints. These effects have been attributed to the irritating nature of the bristles found on the caterpillars. In some instances the reactions have been severe enough to require medical attention (USDA, 1995, vol. III, chapter 3, pp. 2-3), (Allen et al., 1991), (Tuthill, et al., 1984), (Aber, et al., 1982), (Beaucher and Farnham, 1982), (Shama, et al., 1982).

Agricultural, horticultural and forestry enterprises are dependent upon markets beyond the borders of Washington State. Washington must be able to comply with the plant pest and

disease regulations of the Federal government, other states, and international markets. The establishment and spread of the gypsy moth in Washington State would result in the imposition of quarantines (USDA, 1995, vol. II, chapter 2, p. 29). The levels of production and value of plant products would be adversely affected.

2. Proposed Action

a. Bacillus thuringiensis var. (kurstaki) (Btk)

The use of Btk for the eradication of isolated gypsy moth infestations is expected to have no adverse impact on human health or the environment. Various strains of *Bacillus thuringiensis* (B.t.) are a naturally occurring bacterial component of soils worldwide. Modern aqueous formulations of Btk used in gypsy moth control projects contain no organic solvents and have an excellent safety record associated with their use in gypsy moth suppression and eradication projects. An exemption from the requirement of a tolerance has been established for residues of Btk in or on all raw agricultural commodities. This exemption stipulates that manufacturers of Btk test each lot for pathogenicity and vertebrate toxicity. See Appendix D for each Sample Label and MSDS.

A detailed discussion of the human health effects of Btk may be found in the 2004 USDA Forest Service Btk risk assessment. (USDA, 2004)

Due to advances in scientific knowledge, the law requires that pesticides which were first registered before November 1, 1984 be reregistered to ensure that they meet today's more stringent standards. In March of 1998 the United States Environmental Protection Agency came out with a Reregistration Eligibility Decision (EPA, 1998) in which they concluded:

Based on the reviews of the generic data for the active ingredient *Bacillus thuringiensis*, the Agency has sufficient information on the health effects of *Bacillus thuringiensis* and on its potential for causing adverse effects in fish and wildlife and the environment. The Agency has determined that *Bacillus thuringiensis* products, manufactured, labeled and used as specified in this Reregistration Eligibility Decision, will not pose unreasonable risks or adverse effects to humans or the environment. Therefore, the Agency concludes that products containing *Bacillus thuringiensis* for all uses are eligible for reregistration (EPA, 1998).

In the spring of 1999, Foray 48B was applied by aircraft to 52 square miles of Southern Vancouver Island to combat an infestation of European gypsy moth. Approximately 80,000 residents lived in the spray zones. The Capital Health Region coordinated a human health study of possible short-term health effects. The resulting report (Capital Health Region, 1999) concluded:

The results of this project did not show a relationship between aerial spraying of Foray 48B and short-term human health effects. Although some people self-reported health problems that they attributed to the spray program, the research and surveillance methods used in this project did not detect any change in health status

that could be linked to the spray program. Our results showed that many of the health complaints people reported during the spray were as common in people before the spray as they were shortly after the spray. This conclusion is consistent with those of previous studies of the possible health effects of Btk- based pesticide spray programs.

Exposure to Btk spray resulting from its use as proposed in this gypsy moth eradication project is unlikely to cause significant human health effects. However, it is good practice to minimize exposure to any insecticide. One of the conclusions reached in the Oregon study by Green, et al. (1990), was that, "the level of risk for Btk and other existing or future microbial pesticides in immunocompromised hosts deserves further study."

b. Bond Max

Bond Max may be used during ground-based treatments as an adjuvant with the insecticide. Bond Max is a non-ionic spreader-sticker which acts as an adjuvant when mixed with insecticides. Bond Max is not an eye or primary skin irritant per the Federal Hazardous Substances Labeling Act. In the unlikely event that over exposure were to occur, local irritation might be possible, especially in sensitive individuals. Systemic toxic effects are unlikely. See Appendix D for a Sample Label and MSDS.

c. General Precautions

The WSDA will take the following additional steps to assist the public in avoiding or reducing exposure to the spray material:

1. The Pesticide Sensitive Individuals database, maintained by the Pesticide Management Division of the WSDA, will be checked for people living in or near the proposed treatment area who require advance notification.
2. The WSDA will offer a toll-free telephone line with information regarding scheduled treatment days.
3. The WSDA will provide notification calls the day before scheduled applications to any resident in the proposed treatment area requesting them.
4. During ground treatments WSDA on-site spray block monitors will notify residents before the actual application to their property.
5. During ground treatments WSDA on-site spray block monitors will notify bicyclists, joggers and other pedestrians that they are approaching the treatment area.
6. Information will be provided to residents of the treatment area about how to avoid or reduce exposure to the spray material.

B. Non-Target Organisms

1. Bacillus thuringiensis var. (kurstaki) (Btk)

a. Animals

A detailed discussion of the ecological effects of Btk on non-target organisms may be found in the 1995 FEIS vol. II, chapter 4, pp. 52-55, and in vol. IV, chapter 5, pp. 5-10.

As used in gypsy moth eradication projects, Btk has not been shown to adversely affect fish, birds, mammals, or most non-target insects, including honey bees (USDA, 1995, vol. II, chapter 4, pp. 54-55). It is expected that Btk may kill other lepidopteran larvae (leaf-eating caterpillars) if they are present in project areas when treatments occur. In turn, animals dependent on caterpillars as food theoretically may be affected. However, reductions in native caterpillar populations are expected to be temporary due to the brief residual effectiveness of Btk deposits on foliage (4 to 10 days), the high reproductive capacity of most lepidoptera, and recolonization from adjacent untreated areas (USDA, 1995, vol. II, chapter 4, pp. 54-55). The small size of the proposed treatment areas should aid in the recolonization process.

A study conducted in Oregon in connection with gypsy moth control programs in 1986 and 1987 found reduced numbers of caterpillars immediately following Btk treatments and reduced species diversity. This study also found that recovery in numbers of non-target caterpillars began the same season, but that recovery of species diversity lagged behind (Miller, 1990).

One study has shown that Btk could interfere with the biological control of the noxious weed tansy ragwort by cinnabar moth larvae if applied to areas where the weed occurs when late-instar larvae are active (James, et al., 1993). However, an intentionally introduced species of flea beetle has more impact as the primary biological control agent on tansy ragwort (L.C. Burrill, et al. 1994). It is not anticipated that this proposed project would have any adverse impact on flea beetle populations.

Two studies examined the indirect effect of Btk on the reproductive success of insectivorous birds through a possible reduction in food supply. The studies reported no significant differences between treated and untreated areas in numbers of eggs hatched or in nestling growth and development. When caterpillars weren't available, the birds switched to other available prey (Gaddis, 1987), (Gaddis and Corkran, 1986).

There is no evidence of significant adverse impacts of Btk on aquatic organisms. In a study conducted on a benthic stream community there was no evidence that addition of Btk to stream mesocosms created adverse effects for these communities even at greater than 100 times expected exposure rates (Richardson and Perrin, 1994).

b. Plants

Btk is non-toxic to plants. Btk is sensitive to meteorological effects once it has been applied to plant surfaces. Btk is readily removed from plant surfaces by rain and is rapidly degraded by sunlight (USDA, 1995, vol. IV, chapter 7, pp. 15). The use of Bond Max with ground-based equipment will help slow the removal and degradation of Btk by both rain and sunlight.

Changes in soil productivity and fertility due to Btk are not likely. Btk persists for a relatively short time, B.t. is known to occur naturally in soils worldwide, and applications of insecticides containing B.t. do not appear to increase levels of B.t. in soil (USDA, 1995, vol. I, p. 19). For more information about the fate of Btk in the soil refer to 1995 FEIS, vol. 4, chapter 7, p. 16.

c. Threatened, Endangered, and Sensitive Species

No threatened, endangered, or sensitive species are known to be in or near the proposed treatment sites. In reference to the species listed in the Affected Environment section of this EA, all occur well outside of the proposed treatment sites. Therefore, it is not anticipated that the proposed use of Btk would adversely affect these named species.

2. Bond Max

Bond Max may be used during ground-based treatments as an adjuvant with the insecticide. Bond Max is a non-ionic spreader-sticker which acts as an adjuvant when mixed with insecticides. There is no anticipated impact to non-targets.

VI. MONITORING

During the treatment operation, a WSDA or USDA monitor will observe mixing and application of the spray material to ensure compliance with all federal, state, and local laws and regulations and adherence to the Standard Operating Procedures. See Appendix C.

The treatment site will be intensively monitored in the summer of 2011 and 2012 using pheromone-baited traps to determine the effectiveness of the treatment, assist in the eradication and delimit any residual populations of gypsy moths. The results of this monitoring will dictate the need for future action.

VII. CUMULATIVE EFFECTS

The Puyallup site was treated by WSDA with Btk in 2011 and the Eatonville site has not been treated in the past by WSDA.

The most recent Btk application for gypsy moth (prior to 2011) in the Puyallup area occurred in 1995. The most recent Btk treatment in Pierce County occurred in 1998. It is not expected that the proposed 2011 applications will have any cumulative impact on lepidopterans or nontarget organisms.

VIII. SUMMARY

This EA has analyzed the potential environmental effects of the proposed WSDA and USDA APHIS treatment program. This analysis was based on the 1995 USDA FEIS entitled, "Gypsy Moth Management in the United States: a cooperative approach" and the preferred alternative strategy proposed by the Washington State Department of Agriculture and USDA-APHIS for eradicating Gypsy moths at one site in Washington State. The WSDA/USDA-APHIS gypsy moth eradication project strategy proposed for 2011 includes the use of the biological insecticide (Btk) and the spreader-sticker Bond Max during ground-based treatments, followed up by trapping, visual inspections and removal of egg masses where appropriate. It is believed that this IPM strategy will give the project the best chance of achieving the goal of eradicating the gypsy moth infestation while minimizing risks to human health and the environment.

To summarize:

- A. Btk used as described in this Environmental Assessment presents minimal risk of significant impact on human health.
- B. It is not anticipated that any non-target animal or plant populations would be adversely affected due to the limited size of the treatment area. Any detrimental effects on susceptible non-target organisms would be transient and these populations would recover as individuals from nearby untreated areas re-colonized the treatment areas.
- C. No threatened, endangered, or sensitive species would be adversely affected by this eradication project.
- D. No detrimental effects on vegetation, water, or soil are known or anticipated due to this eradication project.
- E. No cumulative effects are known or anticipated.

IX. LIST OF AGENCIES AND PERSONS CONSULTED/NOTIFIED

USDA APHIS Environmental Services; Riverdale, MD

National Marine Fisheries Service for review of the proposed treatment area for the presence of sensitive species or habitats.

US Fish and Wildlife Service for review of the proposed treatment area for the presence of sensitive species or habitats

Washington State Department of Health, Wayne Clifford, for review of the proposed treatment with regard to human health concerns

Tacoma/Pierce County Health Department, Nedda Davies, for review of the proposed treatment with regard to human health concerns

Washington State Department of Natural Resources, Natural Heritage Program, for review of the proposed treatment area for the presence of sensitive species or habitats

Washington State Department of Fish and Wildlife, Ms. Lori Guggenmos, for review of the proposed treatment area for the presence of sensitive species or habitats

Washington State Department of Fish and Wildlife, Ms. Ann Potter, for review of the proposed treatment area for the presence of sensitive lepidopteran species

Washington Department of Ecology for NPDES and SEPA review

Pierce County, Public GIS

City of Puyallup

City of Eatonville

X. LIST OF PREPARERS

Randy Taylor
Gypsy Moth Eradication Coordinator
Washington State Department of Agriculture
3939 Cleveland Ave. SE
Olympia, WA 98501
1-800-443-6684

XI. APPENDICES

- A. References
- B.. Treatment Site Maps
- C. Standard Operating Procedures
- D. Product Labels & Material Safety Data Sheets

APPENDIX A

REFERENCES

Aber, R., DeMelfi, T., Gill, T., Healey, B., Oswell, N., Ruhig, W., Speziale, H., and Witte, E. J. 1982. Rash illness Associated with Gypsy Moth Caterpillars--Pennsylvania. Morbidity and Mortality Weekly Report. 31:13, pp. 169-70.

Allen, V.T.; Gredmiller, O.; Tyler, W.B. 1991. Gypsy Moth Caterpillar Dermatitis Revisited. Journal of the American Academy of Dermatology. 24(6/1): 979-981.

Beaucher, W. N., and Farnham, J.E. 1982. Gypsy-Moth-Caterpillar Dermatitis. The New England Journal of Medicine. 306:21, pp. 1301-02.

Burrill, L.C., Callihan, R.H., Parker, R., Coombs, E., Radtke, H. 1994. Tansy Ragwort Senecio jacobaea L. A Pacific Northwest Extension Publication, Oregon , Idaho, Washington.

Capital Health Region Office of the Medical Health Officer Director of Research. 1999. Human Health Surveillance During the Aerial Spraying for Control of North American Gypsy Moth on Southern Vancouver Island, British Columbia, 1999. December 31, 1999, p 34.

Gaddis, P.K., and Corkran, C.C. 1986. Secondary Effects of B.t. Spray on Avian Predators: the Reproductive Success of Chestnut-backed Chickadees.

Gaddis, P.K. 1987. Secondary Effects of B.t. Spray on Avian Predators: the Reproductive Success of Chickadees.

Green, M., Heumann, M., Sokolow, R., Foster, L.R., Bryant, R., and Skeels, M. 1990. Public Health Implications of the Microbial Pesticide Bacillus thuringiensis: An Epidemiological Study, Oregon, 1985-86. American Journal of Public Health. 80:7, pp. 848-52.

James, R.R., Miller, J.C., and Lighthart, B. 1993. Bacillus thuringiensis var. kurstaki Affects a Beneficial Insect, the Cinnabar Moth (Lepidoptera: Arctiidae). Journal of Economic Entomology. 86:2, pp.334-39.

Miller, J.C. 1990. Field Assessment of the Effects of a Microbial Pest Control Agent on Nontarget Lepidoptera. American Entomologist. 37, pp. 135-39.

Richardson, John S.; Perrin, Chris J. 1994. Effects of the Bacterial Insecticide *Bacillus thuringiensis* var. *kurstaki* (B.t.k.) on a Stream Benthic Community. *Canadian Journal of Aquatic Sciences*. 51.

Shama, S. K., Etkind, P. H., Odell, T. M., Canada, A. T., Finn, A. M., and Soter, N. A. 1982. Gypsy Moth Caterpillar Dermatitis. *The New England Journal of Medicine*. 306:21, pp. 1300-01.

Tuthill, R.W., Canada, A.T., Wilcock, K., Etkind, P.H., Odell, T. M., and Shama, S.K. 1984. An Epidemiologic Study of Gypsy Moth Rash. *American Journal of Public Health*. 74:8, pp. 799-803.

U. S. Department of Agriculture. 1995. Gypsy Moth Management in the United States: a cooperative approach. Final Environmental Impact Statement as Supplemented-- 1995. USDA Animal and Plant Health Inspection Service and USDA Forest Service. Washington D.C.


USDA Forest Service. 2004. Control/Eradication Agents for the Gypsy Moth – Human Health and Ecological Risk Assessment for *Bacillus thuringiensis* var. *kurstaki* (B.t.k.)

U. S. Environmental Protection Agency. 1998. Reregistration Eligibility Decision (RED) *Bacillus thuringiensis*. EPA738-R-98-004. March 1998, pp. 32-33.

APPENDIX B

TREATMENT SITE MAPS

Puyallup Proposed Gypsy Moth Eradication Zone, 2012



South Hill Mall

HI 16

HI 5

39TH

HI 126

HI 112

Legend


Roads

Proposed Eradication Zone

Cartography by: Landon Udo
3/8/2012 Imagery: Bing Maps

0 190 380 760 Feet

Puyallup Proposed Gypsy Moth Eradication Zone Vicinity, 2012



Legend

- Proposed Eradication Zone
- Cities
- Highways
- COUNTY

Cartography by: Landon Udo
3/8/2012

0 30 60 Miles



Puyallup Proposed Gypsy Moth Eradication, Topographic Map 2012



Legend

10 Ft Contours

Proposed Eradication Zone

Cartography by: Landon Udo
3/8/2012 Imagery: Bing Maps

0 230 460 920 Feet

Eatonville Proposed Gypsy Moth Eradication Zone, 2012




Legend

-  Roads
-  Proposed Eradication Zone

Cartography by: Landon Udo
3/8/2012 Imagery: Bing Maps

0 162.5 325 650 Feet

Eatonville Proposed Gypsy Moth Eradication Zone Vicinity, 2012




Legend

- Proposed Eradication Zone
- Cities
- Highways
- COUNTY

Cartography by: Landon Udo
3/8/2012

Eatonville Proposed Gypsy Moth Eradication, Topographic Map 2012



Legend

- 10 Ft Contours
- Proposed Eradication Zone

Cartography by: Landon Udo
3/8/2012 Imagery: Bing Maps

0 180 360 720 Feet

APPENDIX C

WASHINGTON STATE DEPARTMENT OF AGRICULTURE

STANDARD OPERATING PROCEDURES

2012 Gypsy Moth Eradication Project

1. The health and safety of the public, employees of the Washington State Department of Agriculture, and their cooperators will be the first concern in implementing the project.
2. Mixing and application of the insecticide will be done only by an appropriately licensed applicator and will be done only under the supervision of a Washington State Department of Agriculture treatment site monitor.
3. The insecticide will be applied according to label directions.
4. Residents and businesses in the affected eradication area will be notified of the projected dates and times of insecticide applications through email, direct mailings, open house presentations, and press releases. Additionally, a manned 1-800 hotline will be established to address further resident concerns, comments, and project suggestions. Recommendations concerning health and welfare issues will be included in public outreach efforts.
5. The project will commence at the appropriate stage of leaf and/or larval development.
6. Weather conditions, particularly wind, will play the largest role in determining when an effective treatment can be made. In the event of rainfall before spray has had sufficient time to adhere to the foliage, a re-treatment may be necessary.
7. Spill control kits will be on site and readily available during all applications.
8. Treatments will not occur when wind speed exceeds 10 miles/hour.
9. Hydraulic apparatus pressures will be limited to that necessary to obtain thorough coverage to the tops of the tallest trees within the treatment area.

APPENDIX D

**PRODUCT LABELS & MATERIAL SAFETY DATA SHEETS
(MSDS)**

Biological Insecticide

Foray[®] XG

Flowable Concentrate

For Urban, Home and Garden Use

ACTIVE INGREDIENT:

Bacillus thuringiensis, subsp. *kurstaki*, strain
ABTS-351, fermentation solids and solubles 17.19%
OTHER INGREDIENTS 82.81%
TOTAL 100.00%

POTENCY: 10,600 Cabbage Looper Units (CLU/mg) of product (equivalent to 48 billion CLU/GAL).

The % active ingredient does not indicate product performance and potency measurements are not federally standardized.

EPA Reg. No. 73049-46
EPA Est. No. 33762-IA-001

List No. 60178

INDEX:

- 1.0 First Aid
- 2.0 Precautionary Statements
 - 2.1 Hazards to Humans and Domestic Animals
 - 2.2 Personal Protective Equipment (PPE)
 - 2.3 Non-Agricultural Use Requirements
 - 2.4 User Safety Recommendations
 - 2.5 Environmental Hazards
- 3.0 Directions for Use
- 4.0 Non-Agricultural Use Requirements
- 5.0 Application
- 6.0 Mixing
- 7.0 Spray Volumes
- 8.0 General Agricultural Use Instructions
 - 8.1 Application rates
- 9.0 Storage and Disposal
- 10.0 Notice of Warranty

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

1.0

FIRST AID

| | |
|-------------------------------|---|
| If on skin or clothing | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| If in eyes | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-315-9819 (24 hours) for emergency medical treatment and/or transport emergency information. For all other information, call 1-800-323-9597.

2.0 PRECAUTIONARY STATEMENTS

2.1 HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, open wounds or clothing. Wash thoroughly with soap and water after handling.

2.2 Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

2.3 Non-Agricultural Use Requirements:

As a general precaution, when exposed to potentially high concentrations of living microbial products such as this, wear a dust particle mask when mixing or applying this product. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.4 User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.5 Environmental Hazards

Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

3.0 DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CONTINUED

4.0 NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Exposure of unprotected persons can be mitigated by directed spraying. Spray should be allowed to dry undisturbed.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on trees being grown for sale or other commercial use, or for commercial seed production, or for the production of timber or wood products, or for research purposes except wide-area public pest control programs sponsored by government entities, such as mosquito abatement, gypsy moth control, and Mediterranean fruit fly eradication.

Foray XG contains the spores and endotoxin crystals of *Bacillus thuringiensis kurstaki*. Foray XG is a stomach poison and is effective against lepidopterous larvae. After ingestion, larvae stop feeding within hours and die 2-5 days later. Maximum activity is exhibited against early instar larvae. Before use, shake or stir the product. Add some water to the tank mix, pour the required amount of Foray XG into the tank and then add the remaining amount of water to obtain the proper mix ratio. Agitate as necessary to maintain the suspension. Use the diluted mix within 72 hours.

Ground Application

Use an adequate amount of tank mix to obtain thorough coverage without excessive run off. Use the indicated per acre dosages of Foray XG in up to the following amounts of water:

| | |
|--------------------------------|-------------|
| High-volume hydraulic sprayers | 100 gallons |
| Mist blowers | 10 gallons |

5.0 APPLICATION

Foray XG may be applied by ground, undiluted or with quantities of water sufficient to provide thorough coverage of plant parts to be protected. The amount of water needed per acre will depend upon crop size, weather, spray equipment, and local experience.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower/treatment coordinator are responsible for considering all of these factors when making decisions.

6.0 MIXING

Shake or stir Foray XG before use. If dilution is desired, fill spray or mixing tank half of the desired water. Begin agitation and pour Foray XG into water while maintaining continuous agitation. Add other spray material (if any) and balance of water. Agitate as necessary to maintain suspension. Do not allow diluted mixture to remain in the tank for more than 72 hours.

To improve weather-fastness of the spray deposits for hard to wet crops, such as cole crops, use a spreader-sticker approved for use on growing crops. Combinations with commonly used spray tank adjuvants are generally not deleterious to Foray XG, if the mix is used promptly. Before mixing in the spray tank, the testing of physical compatibility by mixing all components in a small container in proportionate quantities will identify possible problems. Checking with an adjuvant supplier for advice on spray adjuvants that are compatible with biological pesticides such as Foray XG, will help avoid incompatibilities.

7.0 SPRAY VOLUMES

Ground Application: Use indicated amount of Foray XG in ground equipment with quantities of water sufficient to provide thorough coverage of plant parts to be protected. The amount of water needed per acre will depend upon crop size, weather conditions, spray equipment used and local experience.

8.0 GENERAL AGRICULTURAL USE INSTRUCTIONS

Foray XG is a biological insecticide for the control of lepidopterous larvae. It contains the spores and endotoxin crystals of *Bacillus thuringiensis kurstaki*. Foray XG must be ingested by the larvae to be effective. For consistent control, apply at first sign of newly hatched larvae (1st and 2nd instar larvae). Susceptible larvae that ingest Foray XG cease feeding within a few hours and die within 2-5 days.

Foray XG may be applied up to and on the day of harvest. For maximum effectiveness follow the instructions listed below:

Monitor to detect early infestations.

Apply Foray XG when eggs start hatching and larvae are small (early instars) and before significant crop damage occurs. Larvae must be actively feeding to be affected.

Repeat applications every 3 to 14 days to maintain control and protect new plant growth. Factors affecting spray interval include rate of plant growth, weather conditions, and reinfestation. Monitor populations of pests and beneficials to determine proper timing of applications.

Under conditions of heavy pest pressures or when large worms are present use the higher rate, shorten the application interval, and/or improve spray coverage to enhance control. When these conditions are present, a contact insecticide can enhance control.

Thorough coverage is essential for optimum performance. Ground applicators equipped with directed drop nozzles can improve coverage.

8.1 Application Rates

| Crop | Pests | Rate ⁽¹⁾ (oz./1000 ft ²) | |
|---|---|--|------------|
| Forests and Shade Trees, Ornamentals, Shrubs, Sugar Maple Trees, Seed Orchards, Ornamental Fruit, Nut and Citrus Trees ⁽²⁾ | Gypsy Moth & Asian Gypsy Moth, Elm Spanworm | 0.5 - 2.5 | |
| | Spruce Budworm, Brown-tail Moth, Douglas Fir Tussock Moth, Coneworm, Buck Moth | 0.5 - 1.9 | |
| Fruiting Vegetables such as: Eggplant, Peppers, Tomatoes | Tussock Moths, Pine Butterfly, Bagworm, Leafrollers, Tortrix, Mimosa Webworm, Tent Caterpillar, Jackpine Budworm, Blackheaded Budworm, Saddled Prominent, Saddleback Caterpillar, Eastern and Western Hemlock Looper, Orangestriped Oakworm, Satin Moth | 0.3 - 1.0 | |
| | Redhumped Caterpillars, Spring and Fall Cankerworm, California Oakworm, Fall Webworm | 0.25 - 0.5 | |
| | Imported Cabbageworm, Diamondback Moth, Green Cloverworm | 0.3 - 0.5 | |
| | Homworms | 0.15 - 1.0 | |
| | Tomato Fruitworm (Heliothis), Variegated Cutworm, Saltmarsh Caterpillar, Loopers | 0.5 - 1.0 | |
| | Armyworms* | 0.5 - 1.8 | |
| | European Corn Borer | 1.0 - 1.3 | |
| | Small Fruit and Berries such as: Blackberries, Blueberries, Currants, Raspberries, Strawberries, Cranberries | Gypsy Moth & Asian Gypsy Moth, Blueberry Leafroller, Loopers, Fruittree Leafroller, Grape Berry Moth, Oblique Banded Leafroller, Achema Sphinx Moth (Hornworm) | 0.5 - 1.0 |
| | | Armyworms* | 0.5 - 1.8 |
| | Brassica (Cole) Vegetables such as: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kohlrabi | Hornworms | 0.15 - 1.0 |
| Webworms, Loopers, Cutworms, Saltmarsh Caterpillar, Omnivorous Leafroller | | 0.5 - 1.0 | |
| Ornamentals, Flowers, Bedding Plants | Diamondback Moth, Imported Cabbageworm, Green Cloverworm | 0.3 - 1.0 | |
| | Armyworms* | 0.5 - 1.8 | |
| | European Corn Borer | 1.0 - 1.3 | |
| | Armyworms* | 0.5 - 1.8 | |
| Greenhouse and Outdoor Nursery Crops such as: Flowers, Brassica, Fruiting Groups, Herbs, and Leafy Vegetables | Azalea Moth, Diamondback Moth, Ello Moth (Hornworm), Io Moth, Loopers, Oleander Moth, Omnivorous Leafroller, Omnivorous Looper, Tobacco Budworm | 0.3 - 0.5 | |
| | Armyworms* | 0.5 - 1.8 | |
| | Heliothis spp, Loopers | 0.3 - 0.5 | |

Special Instructions

* Armyworm Control: Foray XG may be used to control small armyworms (first and second instar) when populations are light and full coverage sprays are applied. Repeat treatment as necessary. If late instar larvae or heavy populations are present, greater control can be achieved by adding a contact insecticide.

(1) Use the higher rates on advanced larval stages or under high density larval populations.

(2) In treating Gypsy Moth and Asian Gypsy Moth infected trees and shrubs in urban, rural, and semi-rural areas, exposure of non-target vegetation including, but not limited to, native and ornamental species and food or feed crops is permitted.

This product can be mixed and used with other pesticides only in accordance with the most restrictive of label limitations and precautions. This product cannot be mixed with any product containing a label prohibition against such mixing. No label dosage rates may be exceeded.

For smaller spray volumes mix the proper number of teaspoons of Foray XG from the following chart to attain the desired rates:

| If the rate is: | Add this amount per gallon of mix: |
|--------------------------------|------------------------------------|
| 0.15 oz./1000 ft. ² | 1/2 teaspoon |
| 0.3 oz./1000 ft. ² | 1 teaspoon |
| 0.5 oz./1000 ft. ² | 1-1/2 teaspoons |
| 1.0 oz./1000 ft. ² | 3 teaspoons |
| 1.3 oz./1000 ft. ² | 4 teaspoons |
| 1.8 oz./1000 ft. ² | 5-1/2 teaspoons |

9.0 STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal of waste.

Pesticide Storage: Store in a cool, dry place. Keep containers tightly closed when not in use. Store in temperatures above freezing and below 32°C (90°F).

Pesticide Disposal: Pesticide waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility in accordance with federal and local regulations.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Home Garden Use Disposal Instructions

Securely wrap original container in several layers of newspaper and discard in trash.

10.0 NOTICE OF WARRANTY

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

FORAY® XGMSDS# BIO-0009C
ISSUED 01/31/05**1. CHEMICAL PRODUCT AND COMPANY IDENTIFICATION**

MATERIAL NAME: Foray® XG
EPA Reg. No.: 73049-46
Code Number: 11046, 12280, 34296
List Number: 60178, 60179, 60180

SYNONYMS:

Biobit® XL
DiPel 48A
Bactospeine XL
Foray 48BA
Foray 48B

MANUFACTURER:

Valent BioSciences Corporation
870 Technology Way, Suite 100
Libertyville, Illinois 60048

EMERGENCY TELEPHONE NUMBERS

Emergency Health or Spill:
Outside the United States: 651-632-6184
Within the United States: 877-315-9819

2. COMPOSITION/INFORMATION ON INGREDIENTS

INGREDIENT NAME: Bacillus thuringiensis, var. kurstaki

CONCENTRATION: 17.19%

CAS NUMBER: 68038-71-1

OSHA-PEL

8HR TWA: N/L

STEL: N/L

CEILING: N/L

ACGIH-TLV

8HR TWA: N/L

STEL: N/L

CEILING: N/L

OTHER LIMITS

8HR TWA: N/A

STEL: N/A

CEILING: N/A

INGREDIENT NAME: Inert/Other ingredients - Proprietary Information

CONCENTRATION: 82.81%

CAS NUMBER: N/A

OSHA-PEL

8HR TWA: N/L

STEL: N/L

CEILING: N/L

ACGIH-TLV

8HR TWA: N/L

STEL: N/L

CEILING: N/L

OTHER LIMITS

8HR TWA: N/A

STEL: N/A

CEILING: N/A

3. HAZARDS INFORMATION

EMERGENCY OVERVIEW: Product is non-toxic by ingestion, skin contact, or inhalation. May be irritating to skin and eyes.

ROUTE(S) OF ENTRY:

Skin: No

Inhalation: No

Ingestion: No

SKIN CONTACT: Mild irritant

SKIN SENSITIZATION: Possible mild sensitizer (unconfirmed)

EYE CONTACT: Mild irritant

TARGET ORGANS: N/D

CARCINOGENICITY RATING:

NTP: N/L

IARC: N/L

OSHA: N/L

ACGIH: N/L

None

SIGNS AND SYMPTOMS: Direct contact with eyes or skin may cause mild irritation.

MEDICAL CONDITIONS AGGRAVATED BY EXPOSURE: N/D

4. FIRST AID MEASURES

EYES: Remove from source of exposure. Flush with copious amounts of water. If irritation persists or signs of toxicity occur, seek medical attention. Provide symptomatic/supportive care as necessary.

SKIN: Remove from source of exposure. Flush with copious amounts of water. If irritation persists or signs of toxicity occur, seek medical attention. Provide symptomatic/supportive care as necessary.

INGESTION: Remove from source of exposure. If signs of toxicity occur, seek medical attention. Provide symptomatic/supportive care as necessary.

INHALATION: Remove from source of exposure. If signs of toxicity occur, seek medical attention. Provide symptomatic/supportive care as necessary.

5. FIRE FIGHTING PROCEDURES

FLASH POINT: N/A (Aqueous suspension)

FLASH POINT METHOD: N/A

LOWER EXPLOSIVE LIMIT(%): N/A

UPPER EXPLOSIVE LIMIT(%): N/A

AUTOIGNITION TEMPERATURE: N/A

FIRE & EXPLOSION HAZARDS: Non-flammable and no explosive properties.

EXTINGUISHING MEDIA: Use appropriate media for underlying cause of fire.

FIRE FIGHTING INSTRUCTIONS: Wear protective clothing and self-contained breathing apparatus.

6. ACCIDENTAL RELEASE MEASURES

SPILL OR RELEASE PROCEDURES: Recover product and place in an appropriate container for disposal. Ventilate and wash the spill area.

7. HANDLING AND STORAGE

HANDLING: The usual precautions for handling chemicals should be observed.

STORAGE: Store in a closed container in a cool, dry place.

SPECIAL PRECAUTIONS: Wash thoroughly with soap and water after handling. Keep impervious gloves on until all potentially contaminated personal protective equipment is removed.

8. EXPOSURE CONTROLS/PERSONAL PROTECTION

ENGINEERING CONTROLS: Use local exhaust

RESPIRATORY PROTECTION: Not usually required. If necessary, use a dust/mist respirator meeting NIOSH standards of at least N-95, R-95 or P-95.

SKIN PROTECTION: Impervious gloves, clothing to minimize skin contact.

EYE PROTECTION: Not usually required. If necessary, use safety glasses or goggles.

OTHER PROTECTION: Wash thoroughly with soap and water after handling.

9. PHYSICAL AND CHEMICAL PROPERTIES

APPEARANCE/PHYSICAL STATE: Light brown aqueous suspension

ODOR: Pungent, musty odor

BOILING POINT: N/D

MELTING/FREEZING POINT: N/D

VAPOR PRESSURE (mm Hg): N/D

VAPOR DENSITY (Air=1): N/D

EVAPORATION RATE: N/D

BULK DENSITY: 1.12-1.2 g/cm³

SPECIFIC GRAVITY: N/D

SOLUBILITY: Readily mixable with water

pH: 4.1-4.8 as a 10% solution in water

VISCOSITY: N/D

10. STABILITY AND REACTIVITY

CHEMICAL STABILITY: Not chemically reactive.

INCOMPATIBILITIES: Alkalinity inactivates product.

HAZARDOUS DECOMPOSITION PRODUCTS: Not known to occur.

HAZARDOUS POLYMERIZATION: Not known to occur.

11. TOXICOLOGICAL INFORMATION**Acute Toxicity**

ORAL LD50: N/D. > 5,000 mg/kg (rat) for a similar formulation. EPA Toxicity Category IV

DERMAL LD50: N/D. > 2,500 mg/kg (rabbit) for a similar formulation. EPA Toxicity Category III

INHALATION LC50: N/D. In a nose-only inhalation study with rats with a similar formulation, no lethality was observed at the highest attainable aerosol concentration of 6.81 mg/liter for 4 hours.

CORROSIVENESS: N/D. Not expected to have any corrosive properties.

DERMAL IRRITATION: Transient, slight or mild irritation noted in a dermal irritation study with a similar formulation. EPA Toxicity Category IV.

OCULAR IRRITATION: Transient, mild irritation was observed in test animals in a study a similar formulation. EPA Toxicity Category III.

DERMAL SENSITIZATION: N/D. The possibility of mild sensitization exists with this formulation, however, this has not been confirmed by actual experience.

SPECIAL TARGET ORGAN EFFECTS: N/D

CARCINOGENICITY INFORMATION: N/D. None of the components are classified as carcinogens.

12. ECOLOGICAL INFORMATION

ECOLOGICAL INFORMATION: Studies on non-targets have been performed without identifying any organisms at risk. The following species have been included in the testing: mammals (rats, rabbits); freshwater aquatic organisms (Daphnia magna, Rainbow Trout); birds (Mallard, Bobwhite); and non-target insects (Green Lacewing larvae, Ladybird Beetles, Honey Bee).

13. DISPOSAL CONSIDERATIONS

WASTE DISPOSAL METHODS: Dispose of product in accordance with federal, state and local regulations.

14. TRANSPORTATION INFORMATION**DOT**

STATUS: Not Regulated

PROPER SHIPPING NAME: N/A

HAZARD CLASS: N/A

UN NUMBER: N/A

PACKING GROUP: N/A

REPORTABLE QUANTITY: N/A

IATA/ICAO

STATUS: Not Regulated

PROPER SHIPPING NAME: N/A

HAZARD CLASS: N/A

UN NUMBER: N/A

PACKING GROUP: N/A

REPORTABLE QUANTITY: N/A

IMO

STATUS: Not Regulated

PROPER SHIPPING NAME: N/A

HAZARD CLASS: N/A

UN NUMBER: N/A

PACKING GROUP: N/A

REPORTABLE QUANTITY: N/A

FLASH POINT: N/A

15. REGULATORY INFORMATION

TSCA STATUS: Exempt

CERCLA STATUS: N/D

SARA STATUS: N/D

RCRA STATUS: N/D

PROP 65 (CA): N/D

16. OTHER INFORMATION

REASON FOR ISSUE: Added alternate brand name (synonym) - Foray XG

APPROVAL DATE: 07/20/04

SUPERSEDES DATE: 06/11/04

Note: Combined and Replaced MSDS # BIO-0033 Rev 0.

LEGEND:

N/A = Not Applicable

N/D = Not Determined

N/L = Not Listed

L = Listed

C = Ceiling

S = Short-term

® = Registered Trademark of Valent BioSciences

™ = Registered Trademark of Valent BioSciences
The information and recommendations contained herein are based upon tests believed to be reliable. However, Valent BioSciences does not guarantee their accuracy or completeness nor shall any of this information constitute a warranty, whether expressed or implied, as to the safety of the goods, the merchantability of the goods, or the fitness of the goods for a particular purpose. Adjustment to conform with actual conditions of usage may be required. Valent BioSciences assumes no responsibility for results obtained or for incidental or consequential damages arising from the use of these data. No freedom from infringement of any patent, copyright or trademark is to be inferred.

VALENT BIOSCIENCES™ CORPORATION
870 Technology Way, Suite 100
Libertyville, IL 60048 - 800-323-9597
July 2004

© Valent BioSciences Corporation VID 1.31.05

BOND[®] MAX

SPREADER STICKER DEPOSITION AID

Principal Functioning Agents:

Alcohol ethoxylate, 1,2-propanediol and synthetic latex57.5%
 Constituents ineffective as spray adjuvant42.5%
TOTAL100.0%

CA Reg. No. 34704-50060
 WA Reg. No. 34704-08903

**KEEP OUT OF REACH OF CHILDREN
 WARNING**

NET CONTENTS: 2.5 U.S. GALLONS (9.462 L)



Loveland Products, Inc. • PO Box 1286 • Greeley, CO 80632-1286

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash clothing before reuse. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.
Personal Protective Equipment: Wear protective eyewear such as goggles, face shield, or safety glasses and wear long-sleeved shirt and long pants, socks, shoes and gloves.

First Aid: If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL 1-866-944-8565.

GENERAL: BOND MAX[®] combines the adherence ability of a latex polymer with super coverage of non-ionic surfactants. BOND MAX will increase deposition, reduce run-off and secure spray from rain or overhead irrigation. BOND MAX will decrease surface tension, resulting in better coverage. BOND MAX promotes rapid spreading for a uniform protective film. Apply sprays containing BOND MAX at least 30 minutes before an anticipated rain or overhead irrigation.

DIRECTIONS FOR USE: Fill spray tank 1/2 full with water and begin agitation. Add pesticides as directed by label, while maintaining agitation. After pesticides are thoroughly mixed, eliminate any existing foam with approved defoamer (Unifoamer[®]). Then fill tank to desired water volume before adding BOND MAX. For tank mix compatibility concerns, conduct a jar test of the proposed mixture to ensure compatibility of all components. Mix components in the same ratio as the proposed tank mix.

Suggested Use Rates: Field & Row Crops: Ground and Air Application (fl. oz./100 gallons):

Above 50 GPA: 12 to 24
 10 to 50 GPA: 20 to 32
 Below 10 GPA: 28 to 40

| Gallons of Spray Volume per Acre | | BOND MAX per 100 gallons spray |
|----------------------------------|-------------------|--------------------------------|
| Vineyards, hop yards | Treefruits & nuts | |
| | 150 to 300 | 1 to 2 ounces |
| | 75 to 150 | 2 to 4 ounces |
| | 50 to 100 | 3 to 6 ounces |
| | 20 to 40 | 8 to 16 ounces |
| | 15 to 30 | 10 to 20 ounces |
| | 8 to 16 | 1 pint to 1 quart |
| | 5 to 10 | 1.5 pints to 1.5 quarts |

Rinse tank, pump, lines and nozzles with water immediately after spraying. Observe the pre-harvest interval on the pesticide label when using BOND MAX. No time limitations apply to non-food crops.

Environmental Hazards: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash-waters.

STORAGE AND DISPOSAL

STORAGE: Store in cool, dry place. Store in original container. Keep container tightly closed. Do not reuse empty container. **DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes may be disposed of on-site or at an approved waste disposal facility. Triple rinse (or equivalent) adding rinse water to spray tank. Offer container for recycling or dispose of container in sanitary landfill, or by other procedures approved by appropriate authorities. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state and local ACRC recycler visit the ACRC web page at www.acrcycle.org. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary. LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when the product is used in accordance with such Directions for Use under normal conditions of use. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT SHALL BE LIMITED TO ONE OF THE FOLLOWING: AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT; LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES; OR DAMAGES IN THE NATURE OF A PENALTY.

BOND MAX[®] and UNFOAMER[®] are registered trademarks of Loveland Products, Inc.

BOND MAX 2.5GL/A1012

MATERIAL SAFETY DATA SHEET

BOND MAX®

FOR CHEMICAL EMERGENCY, SPILL, LEAK, FIRE, EXPOSURE OR ACCIDENT, CALL CHEMTREC - DAY OR NIGHT 1-800-424-9300

1. CHEMICAL PRODUCT AND COMPANY IDENTIFICATION

FORMULATED FOR:

LOVELAND PRODUCTS, INC.
P.O. Box 1286 • Greeley, CO 80632-1286

24-Hour Emergency Phone: 1-800-424-9300
Medical Emergencies: 1-866-944-8565
U.S. Coast Guard National Response Center: 1-800-424-8802

PRODUCT NAME: BOND MAX® SPREADER STICKER DEPOSITION AID
CHEMICAL NAME: Synthetic Latex (combination of synthetic latex, 1,2-Propanediol and alcohol ethoxylate)
CHEMICAL FAMILY: Mixture
CA REG. NO.: 34704-50060
WA REG. NO.: 34704-08003
MSDS Number: 8111550-10-LPI MSDS Revisions: Sections 1 and 13 Date of Issue: 07/08/10 Supersedes: 10/14/08

2. HAZARDS IDENTIFICATION SUMMARY

KEEP OUT OF REACH OF CHILDREN - WARNING - Causes substantial but temporary eye irritation. Do not get in eyes or on clothing. Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash clothing before reuse. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Primary routes of entry are eye contact and skin contact

This product is a sticking agent with surfactant. This product is a light blue liquid with mild odor.

3. COMPOSITION, INFORMATION ON INGREDIENTS

| <u>Chemical Ingredients:</u> | <u>Percentage by Weight:</u> | <u>CAS No.</u> | <u>TLV (Units)</u> |
|--|------------------------------|----------------|--------------------|
| Synthetic Latex, 1,2-Propanediol, alcohol ethoxylate | 57.50 | Mixture | Not established |
| Other Ingredients | 42.50 | | |

4. FIRST AID MEASURES

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

5. FIRE FIGHTING MEASURES

FLASH POINT (°F/Test Method): >212°F / >100°C (PMCC)
FLAMMABLE LIMITS (LFL & UFL): Not established
EXTINGUISHING MEDIA: Dry chemical or carbon dioxide (CO₂), foam or water spray/fog.
HAZARDOUS COMBUSTION PRODUCTS: Carbon monoxide and/or carbon dioxide
SPECIAL FIRE FIGHTING PROCEDURES: Wear self-contained breathing apparatus and full protective gear.
UNUSUAL FIRE AND EXPLOSION HAZARDS: None.

6. ACCIDENTAL RELEASE MEASURES

STEPS TO BE TAKEN IF MATERIAL IS RELEASED OR SPILLED:

Wear appropriate personal protective equipment (refer to Section 8). Pick up the material with absorbent material and place in a container for proper disposal in accordance with local, state and federal regulations.

ENVIRONMENTAL PRECAUTIONS: Keep spills and cleaning runoff out of municipal sewers and open bodies of water.

7. HANDLING AND STORAGE

HANDLING: Personal Protective Equipment: Wear protective eyewear, such as goggles, face shield, or safety glasses and long-sleeved shirt and long pants, socks, shoes and gloves. Wash thoroughly after handling.
STORAGE: Store in a cool, dry place. Store in original container. Keep tightly closed. Do not reuse empty container. Do not contaminate water, food or feed by storage or disposal.

8. EXPOSURE CONTROLS / PERSONAL PROTECTION

ENGINEERING CONTROLS: Work in well-ventilated area. Local exhaust may be required if working in confined space.
RESPIRATORY PROTECTION: Wear a NIOSH approved air-purifying respirator for pesticide handling if necessary.
EYE PROTECTION: Chemical goggles or face-shield.
SKIN PROTECTION: Wear long sleeved shirt, long pants, shoes and socks.

| | | |
|-----------------|---------------------------------|--|
| 1,2-Propanediol | OSHA PEL 8 hr TWA not listed | AIHA WEELS TWA 10 mg/m ³ |
|-----------------|---------------------------------|--|

9. PHYSICAL AND CHEMICAL PROPERTIES

APPEARANCE AND ODOR: Light blue liquid with mild odor. **SOLUBILITY:** Dispersible
SPECIFIC GRAVITY (Water = 1): 1.026 g/ml **BULK DENSITY:** 8.56 lbs/gal. **pH:** 7.4 (1% solution)
VAPOR PRESSURE: Not established **BOILING POINT:** Not established
PERCENT VOLATILE (by volume): Not established **EVAPORATION RATE:** Not established
 Note: These physical data are typical values based on material tested but may vary from sample to sample.
 Typical values should not be construed as a guaranteed analysis of any specific lot or as specification items.

10. STABILITY AND REACTIVITY

STABILITY: Stable
CONDITIONS TO AVOID: None known.
INCOMPATIBILITY: Low pH (strong acidic conditions) will cause coagulation. Excessive free metallic ions may cause coagulation.
HAZARDOUS DECOMPOSITION PRODUCTS: Carbon monoxide from burning.
HAZARDOUS POLYMERIZATION: Will not occur.

11. TOXICOLOGICAL INFORMATION

Acute Oral LD₅₀ (rat): > 5000 mg/kg **Acute Dermal LD₅₀ (rabbit):** >2000 mg/kg
Eye Irritation (rabbit): Moderate eye irritant **Skin Irritation (rabbit):** Slight skin irritant
Inhalation LC₅₀ (rat): Not established **Skin Sensitization:** Not a sensitizer.
Carcinogenic Potential: Not listed by OSHA, NTP, IARC, and ACGIH as a known human carcinogen

12. ECOLOGICAL INFORMATION

Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

13. DISPOSAL CONSIDERATIONS

Wastes may be disposed of on site or at an approved waste disposal facility. Triple rinse (or equivalent), adding rinse water to spray tank. Offer container for recycling or dispose of in a sanitary landfill or by other procedures approved by appropriate authorities. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state and local ACRC recycler visit the ACRC web page at <http://www.acrcycle.org/>. Do not contaminate water, food or feed by storage or disposal.

14. TRANSPORT INFORMATION

DOT Shipping Description: NOT REGULATED BY USDOT.
Freight Classification: ADHESIVES, ADJUVANTS, SPREADERS OR STICKERS (NMFC 4610; CLASS: LTL 60, TL 35)
 Consult appropriate ICAO/IATA and IMDG regulations for shipment requirements in the Air and Maritime shipping modes.

15. REGULATORY INFORMATION

| | | | |
|--|----------------|------------|----------------|
| NFPA & HMIS Hazard Ratings: | NFPA | | HMIS |
| | 2 Health | 0 Least | 2 Health |
| | 2 Flammability | 1 Slight | 2 Flammability |
| | 0 Instability | 2 Moderate | 0 Reactivity |
| | | 3 High | H PPE |
| | | 4 Severe | |

MATERIAL SAFETY DATA SHEET

BOND MAX®

SARA Hazard Notification/Reporting

SARA Title III Hazard Category:

Immediate Y
Delayed N

Fire N
Reactive N

Sudden Release of Pressure N

Reportable Quantity (RQ) under U.S. CERCLA: Not listed

SARA, Title III, Section 313: Not listed

RCRA Waste Code: Not listed

CA Proposition 65: Not listed

16. OTHER INFORMATION

MSDS STATUS: Sections 1 and 13 revised.

PREPARED BY: Registrations and Regulatory Affairs

REVIEWED BY: Environmental/Regulatory Services

©Bond Max is a registered trademark of Loveland Products, Inc.

Disclaimer and Limitation of Liability: This data sheet was developed from information on the constituent materials identified herein and does not relate to the use of such materials in combination with any other material or process. No warranty is expressed or implied with respect to the completeness or ongoing accuracy of the information contained in this data sheet, and LOVELAND PRODUCTS, Inc. disclaims all liability for reliance on such information. This data sheet is not a guarantee of safety. Users are responsible for ensuring that they have all current information necessary to safely use the product described by this data sheet for their specific purpose.



APPENDIX E

Finding of No Significant Impact

Finding of No Significant Impact
for
PIERCE, COUNTY
WASHINGTON

2012 APHIS Cooperative Gypsy Moth Eradication Program
Site-Specific Environmental Assessment

The United States Department of Agriculture, (USDA), in cooperation with the Washington State Department of Agriculture, (WSDA), proposes an eradication program to eliminate two isolated infestations and/or introductions of the non-native Gypsy Moth, *Lymantria dispar*, (Linnaeus), in Pierce county, Washington during the spring of 2012. Under the process described in the National Environment Policy Act, 1969 (NEPA), an Environmental Assessment (EA) was prepared to analyze the effect of the proposed action at the site-specific level. The environmental consequences of this program are analyzed in this EA, which is supported by and tiered to the "Gypsy Moth Management in the United States: a cooperative approach, Final Environmental Impact Statement, November 1995", (FEIS). The USDA examined the six alternatives available in the FEIS and has selected the preferred Alternative 6, which consists of suppression, eradication, and slow the spread. Under alternative 6, several treatment options are available for Gypsy Moth management. The treatment options analyzed included:

- 1) No action
- 2) *Bacillus thuringiensis* var. *kurstaki* (Btk); a biological insecticide
- 3) Diflubenzuron; a chemical insecticide
- 4) Gypsy Moth nucleopolyhedrosis virus (NPV) or Gypchek; a biological insecticide
- 5) Mass trapping, Gypsy Moth traps with disparlure to attract male Gypsy Moths
- 6) Mating disruption, aerial application of disparlure
- 7) Sterile insect release, release of sterile or partially-sterile Gypsy Moth life stages

The potential environment impacts and mitigation measures of these treatment options are described in the FEIS and EA. The EA was prepared by the USDA and WSDA. The FEIS and EA are available for review at the following locations:

USDA-APHIS-PPQ
Office of the State Plant Health Director
33400 9th Ave. S., Suite 200
Federal Way, WA 98003

Washington State Library
6880 Capitol Blvd. S
Tumwater, WA 98501

USDA-APHIS-PPQ
APHIS Library, 1st floor
4700 River Road
Riverdale, MD 20737


A cooperative USDA/WSDA eradication project is selected. This cooperative program selects the preferred Alternative 6: specifically eradication, due to the geographic location of Washington State. The USDA / WSDA Gypsy Moth eradication strategy proposed for 2012 includes utilizing three to five ground applications of the biological insecticide, *Bacillus thuringiensis* var. *kurstaki*, (*Btk*), applied to all foliage within the treatment areas. Treatments will begin in late April and will conclude by early June. The exact number and timing of treatments will depend larval development, foliage development and weather conditions. The insecticide may be mixed with a spreader-sticker (Bond Max®) during ground treatments. The success of the applications will be monitored by intensive trapping in the summer of 2012.

No comments were received during the 30-day public comment period. For more information on implementation of this program, please refer to the site specific 2012 EA. Implementation of this program, with associated operating procedures and mitigation measures as identified in the EA, would ensure that no significant adverse environmental impact would occur to the human environment.

Reasons for the finding of no significant impact include:

- A. B.t.k. used as described in the Environmental Assessment presents minimal risk of significant impact on human health.
- B. It is not anticipated that any non-target animal or plant populations would be adversely affected due to the limited size of the treatment areas. Any detrimental effects on susceptible non-target organisms would be transient and these populations would recover as individuals from nearby untreated areas re-colonized the treatment areas.
- C. No threatened, endangered, or sensitive species would be adversely affected by this eradication project.
- D. No detrimental effects on vegetation, water, or soil are known or anticipated due to this eradication project.
- E. No cumulative effects are known or anticipated.

This EA is consistent with Executive Order No. 12898, "Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations." That implementation of this cooperative USDA/WSDA eradication project will not result in disproportionately high and adverse human health or environmental effects on any minority populations and low-income populations. As required by the Executive Order of the President, opportunities for full participation in the NEPA process by such populations have been provided.


Barbara A. Chambers

State Plant Health Director – WA State
United States Department of Agriculture
Animal & Plant Health Inspection Service
Plant Protection and Quarantine

04/17/12

Date